

Influence of environmental factors on age at first calving in Jaffarabadi buffaloes

H.A. Sharma^{1*}, P.U. Gajbhiye², A.R. Ahlawat¹, A.L. Ramani¹,
V.B. Dongre³

¹Department of Animal Genetics & breeding, College of Veterinary Science ²Cattle breeding farm, Junagadh Agricultural University, Junagadh, Gujarat, India; ³Cattle Breeding Farm, College of Veterinary Science & A. H., Udgir Maharashtra, India
Corresponding author, Email: hardiksharma.vet@gmail.com

Journal of Livestock Science (ISSN online 2277-6214) 7: 256-259

Received on 12/09/2016; Accepted on 25/9/2016

Abstract

The breeding records of 191 primipara Jaffarabadi buffaloes spread over 1991 to 2014 from Cattle Breeding farm, Junagadh were used to study the effects of periods and seasons of conception on age at first calving. The data were collected and analysed using maximum likelihood and least square techniques to examine the effect of non-genetic factors on age at first calving. The overall least squares mean for Age at first calving was 1788.52 ± 36.14 days. The effect of season of conception was significant ($p < 0.05$) and effect of season of calving and period of calving were highly significant ($p < 0.01$) on age at first calving. Therefore, the data for genetic analysis should be adjusted for better estimates of genetic parameters in Jaffarabadi buffaloes.

Key words: age at first calving; Period of calving, calving season; buffalo; Jaffarabadi

Introduction

India ranks first in Livestock population in the world with 512.05 million heads. Buffaloes are regarded as a good option to dairy breeding, because of their potential milk production in distinct environmental conditions and market value of its milk. India is the treasure house of world's best buffalo germplasm with the population of about 108.7 million in 2012, which was about 57.3 percent of total buffalo population of the world, (DAHDF, 2013). India has continued to be the largest milk producing country in 2014-15 with an anticipated milk production of 146.3 million tonnes. The country's share in world milk production stands at 18.5 percent and buffaloes contribute 51 percent of total milk yield of the country, even though they are less in number than cattle (DAHDF, 2016). About 63 % of the world's buffalo milk and 95% of buffalo milk in Asia is contributed by Indian buffaloes (FAO, 2012).

Jaffarabadi is considered to be one of the best dairy buffalo breed in India. They are heaviest and massive type of riverine buffalo. They are good milkers and thrive well on natural grazing due to their greater feed conversion efficiency. The native breeding tract of Jaffarabadi buffalo is Saurashtra region of Gujarat, viz. Junagadh, Bhavanagar, Jamnagar, Amreli, Gir Somnath, Rajkot and Morbi district as well as some part of Surendranagar district. It is also known as "Bhavnagri", "Gir" or "Jaffari" by the local people (Kathiravan *et al.*, 2007).

The age at first calving (AFC) is an important parameter in reproductive efficiency. Age at first calving is a controversial point not merely among the farmers but also among many animal breeders as they feel that too early age at first calving, has an adverse effect on the future performance of animals (Rakshe, 2003). Even though cattle occupy an important place in the agricultural economy of India, so far as milk production is concerned, the buffalo has taken her place as a milk producing animal. Raising heifers is the most expensive component of the dairy farm operations. Poor growth rate resulting in delayed age at maturity in our local dairy animals further aggravates the situation. However, cost of heifer production can be reduced through better management, balanced feeding, use of performance modifiers and better health care. The shorter the age at first calving the longer shall be productive life (Bhatti *et al.*, 2007).

Considering above background, present study was intended to evaluate effects of various non-genetic factors on age at first calving in Jaffarabadi buffaloes, which will help to formulate suitable evaluation procedures as it is expected that animals with early sexual maturity in terms of AFC may also initiate physiological functioning of reproduction and milk production earlier.

Material and methods

The records on 191 primipara Jaffarabadi buffaloes maintained at cattle breeding farm, Junagadh Agricultural University, Junagadh spread over a period from 1991 to 2014 were analysed to study the effects on non-genetic factors on age at first calving in Jaffarabadi buffaloes.

The collected data were classified and coded into 5 different periods: Period I (1991-1995), Period II (1996-2000), Period III (2001-2005), Period IV (2006-2010) and Period V (2011-2014) due to change in different conditions of climatic cycles, fodder and feed availability and management practices over time. Each year was divided into 5 different seasons: rainy, autumn, winter, spring, summer in order to assess the effect of non-genetic factors on age at first calving.

The influence of various non genetic factors on age at first calving were studied using least square analysis technique described by Harvey (1986). The statistical model used for age at first calving is given below:

$$Y_{ijkl} = \mu + a_i + b_j + c_k + e_{ijkl}$$

Where,

Y_{ijkl} = Observation on the l^{th} individual in i^{th} season, j^{th} period

μ = Overall population mean

a_i = Effect of i^{th} season of calving ($i = 1$ to 5)

b_j = Effect of j^{th} period of calving ($j = 1$ to 5)

c_l = Effect of season of conception ($l = 1$ to 5)

e_{ijk} = Random error, NID ($0, \sigma_e^2$)

Duncan's multiple range test as modified by Kramer (1957) was used for testing significant differences among least squares means.

Results and discussion

The age at first calving is an important economical trait regarding reproduction. The least-squares means for AFC in different seasons of calving and periods of calving are detailed in Table 1. The overall means for AFC was 1788.52 ± 36.14 days in Jaffarabadi buffalo (Table 1).

Table1: Least squares means of age at first calving in Jaffrabadi Buffaloes

Effect	n	Mean \pm SE (in days)
Overall	191	1788.52 \pm 36.14
Season of conception		
Summer	8	1502.03182 \pm 137.62 ^a
Rainy	55	1849.391059 \pm 83.10 ^c
Autumn	81	1901.60276 \pm 81.84 ^c
Winter	30	1934.024179 \pm 89.15 ^c
Spring	17	1763.471483 \pm 129.27 ^b
Season of calving		
Summer	6	2025.48 \pm 121.02 ^d
Rainy	119	1671.45 \pm 30.29 ^a
Autumn	37	1855.68 \pm 49.35 ^c
Winter	21	1762.07 \pm 65.19 ^b
Spring	8	1627.93 \pm 105.65 ^a
Period of calving		
1991 – 1995	34	1631.92 \pm 59.97 ^a
1996 – 2000	17	2023.80 \pm 72.18 ^c
2001 – 2005	37	1780.68 \pm 58.50 ^b
2006 – 2010	55	1838.71 \pm 52.53 ^b
2011 – 2014	48	1667.51 \pm 53.28 ^a

Effect of season of conception on AFC

The season of conception had a significant effect on age at first calving in this study. The present findings were in close agreement with Penchev *et al.* (2014). The significant influence might be due to a distinct seasonal variation in display of oestrus, conception rate in buffaloes having better reproductive efficiency during winter compared to summer months attributed to environmental factors as also documented by Madan (1988) and Tailor (1990).

Effect of season of calving on AFC

The season of calving had a highly significant ($P < 0.01$) effect on age at first calving. The animals calved during summer season showed significantly higher AFC, whereas less in rainy and spring seasons. The buffaloes that calved during hot season showed delayed sexual maturity as compare to cooler seasons because Jaffarabadi buffaloes are partially seasonal breeder. Similar finding were reported by other workers in Murrah buffaloes (Dutt *et al.*, 2001 and Jamuna *et al.*, 2015). Kanaujia *et al.* (1974) also reported highly significant effects of season of year on age at first calving. However, differences between seasons were non-significant for AFC were reported by several workers (Jain and Taneja, 1982, Johari and Bhatt, 1979 and Reddy and Mishra, 1980).

Effect of period of calving on AFC

Period of birth had a statistically significant ($P < 0.01$) effect on AFC. The significant influences indicated that change in population structure, climatic conditions, fodder and feed availability and management practices over time brought variation in AFC. The overlook of table 1 did not show any specific trend in AFC. However, during last period the reduced AFC could be the result of a better management practices over time. The findings in the present study were in agreement with Johari and Bhatt (1979). Charlini and Sinniah (2015) also reported that the age at first calving was influenced by breed and year of calving. However, Jamuna *et al.* (2015) reported a non-significant effect of period on age at first calving.

Conclusion

The age at first calving in Jaffarabadi buffaloes was significantly influenced by period and season of calving. The difference in AFC over different period and season may be attributed to differences in feeding, management practices and differential culling level. Therefore, non-genetic variation should be taken into consideration when developing models to be used in adjusting data to provide best estimates of genetic parameters in Jaffarabadi buffaloes.

References

- 1) Bhatti S.S., Sarwar M., Khan M.S., and Hussain S.M.I. 2007. Reducing the age at first calving through nutritional Manipulations in dairy buffaloes and cows: a review. *Pakistan Veterinary Journal*, **27(1)**: 42–4.
- 2) Charlini B.C. and Sinniah J. 2015. Performance of Murrah, Surti, Nili-Ravi buffaloes and their crosses in the intermediate zone of Sri Lanka. *Livestock Research for Rural Development*, **27(3)**. Article 47. (<http://www.lrrd.org/lrrd27/3/char27047.html>)
- 3) DAHDF. 2013. Annual Report 2012-13. Department of Animal Husbandry, Dairying and Fisheries. Government of India.
- 4) DAHDF. 2016. Annual Report 2015-16. Department of Animal Husbandry, Dairying and Fisheries. Government of India.
- 5) Dutt T., Bhushan B. and Kumar S. 2001. First lactation and lifetime performance traits in Murrah buffaloes. *The Indian journal of Animal Sciences*, **71(5)**: 483– 484.
- 6) FAO. 2012. Buffalo production and research. Food and Agriculture Organization. (<http://www.fao.org/statistics/en>).
- 7) Harvey W.R. 1987. Mixed model least-squares and maximum likelihood computer program. PC-I. Ohio State University, Columbus, U.S.A. (Mimeograph).
- 8) Jain A. and Taneja V.K. 1982. Effect of non-genetic factors on reproduction and production traits in Murrah buffaloes. *Asian Journal of Dairy Research*, **1**: 123–129.
- 9) Jamuna V., Chakravarty A.K. and Patil C.S. 2015. Influence of non-genetic factors on performance traits in Murrah buffaloes. *Indian Journal of Animal Research*, **49(3)**: 279–283.
- 10) Johari D.C. and Bhatt P.N. 1979. Effect of genetic and non-genetic factors on production traits in buffaloes. *The Indian journal of Animal Sciences*, **49**: 984–991.
- 11) Kanaujia A.S., Balaine D.S. and Rathi S.S. 1974. Factors affecting some economic traits of reproduction in Indian buffaloes. *The Indian journal of Dairy science*, **27**: 264–270.
- 12) Kathiravan P., Mishra B.P., Sadana D.K., Katariya, R.S. and Singh, G. 2007. Buffalo Genetic Resources of India: Jaffarabadi. Monograph, National Bureau of Animal Genetic Resources, Karnal.
- 13) Kramer C.Y. 1957. Extension of multiple range tests to group correlated adjusted means. *Biometrics*, **13(1)**: 13–18.
- 14) Madan M.L. 1988. Status of reproduction in female buffalo. In: Buffalo Production and Health: a compendium of latest research information based on Indian studies, ICAR Publication, New Delhi, India, pp. 89-100.
- 15) Penchev P., Ilieva Y., Dimov K. 2014. Effect of season of birth on season of calving and age at first calving in buffalo heifers. *Bulgarian Journal of Agricultural Science*, **20(2)**: 447-451.
- 16) Rakshe P.T. 2003. Effect of age at first calving and subsequent period of breeding on the performance of buffaloes from the college of agriculture Pune (MS), India. *Buffalo Bulletin*, **22(1)**: 7-11.
- 17) Reddy K.M. and Mishra R.R. 1980. Effect of Age at first calving on lactation components and calving distribution in Murrah buffaloes. *The Indian journal of Dairy science*, **33(3)**: 347–50.
- 18) Tailor S.P., Jain, L.S., Gupta, H.K., Bhatia, J.S. 1990. Oestrus and conception rates in buffaloes under village conditions. *The Indian journal of Animal Sciences*, **60**: 1020-1021.